



EGYPTIAN ACADEMIC JOURNAL OF
BIOLOGICAL SCIENCES
ENTOMOLOGY


A



ISSN
1687-8809

WWW.EAJBS.EG.NET

Vol. 16 No. 2 (2023)



Latent Histopathological Effects of Lufenuron and Rice Bran Extract on *Schistocerca Gregaria* (Forsk.) (Orthoptera: Acrididae)

Noura M. Mahdy¹, Samira M. N. Abd-El Wahed¹ and Amira E. Abdel hamid^{2*}

¹Plant Protection Research Institute, Agricultural Research Center, Giza, Egypt.

²Department of Entomology, Faculty of Science, Ain Shams University, Cairo, Egypt

*E-mail: amiraelsayed73@yahoo.com

ARTICLE INFO

Article History

Received:15/4/2023

Accepted:16/6/2023

Available:20/6/2023

Keywords:

Schistocerca gregaria;
histological study;
lufenuron; chitin
synthesis inhibitors;
rice bran extract;
agricultural waste
products.

ABSTRACT

Laboratory experiments evaluated the latent effects of lufenuron and rice bran extract on the newly hatched nymphs of *Schistocerca gregaria* produced from females who were treated as fifth nymphal instar with LC₅₀ of each lufenuron and rice bran extract. Semithin sections of the newly hatchling showed great histological malformations in the structure of the brain, compound eyes, midgut, and hind leg muscles compared with control, these organs lost their normal structures, shapes, and functions. The tested compounds had serious extended histopathological effects, so they can be used to control *S. gregaria* as safe alternative agents to toxic synthetic pesticides.

INTRODUCTION

The main source of food for people and animals, cultivated crops, is severely damaged by the desert locust *Schistocerca gregaria*. Due to its two to five annual generations and the capacity of its swarms to travel quickly over huge distances, the desert locust is arguably the most deadly of the locust pests (Homberg, 2015). In a gregarious phase, the desert locust can result in up to 100% crop loss (Simpson *et al.*, 1999). So, it seemed necessary to develop an effective preventive control strategy against its nymphal instar to stop the emergence of mobile swarms (Taha & El-Gammal, 1990). The main pesticides used to control locusts today are organophosphorus pesticides because of the banning of organochlorines (Lecoq, 2001); the current use of conventional pesticides in desert locust control may have direct and indirect adverse effects on human health and the environment (Garriga, & Caballero, 2011). To solve these problems, alternative agents for controlling desert locusts must be found that are economical, practical, and safe for the environment. In contrast to the dangers of toxic synthetic pesticides, which act as potential acute or chronic insecticides, insect growth regulators (IGRs) and botanical extracts have low mammalian toxicity and low environmental problems.

Chitin synthesis inhibitors are part of insect growth regulators, that effectively inhibit insect development throughout their entire life cycle (Verloop & Ferrel, 1977), the desert locust is effectively killed by these compounds (Azam & Seegh, 1993; Coppen & Jepson 1996; and Bakr *et al.*, 2008). Numerous studies have proven the activity of lufenuron

(CSI) as an inhibitor of ovicidal, transovarial-ovicidal, and embryonic development of pests at the recommended dose (Fonseca *et al.*, 2015 and Sampson *et al.*, 2017). Also, it causes different morphological, biochemical, and histological malformation in different stages of the desert locust (Abdel Rahman, 2017; Al-Zeeb *et al.*, 2018; Mahdy *et al.*, 2019 & 2020 and Said, 2020). As well it has reasonable persistence and low toxicity against non-target organisms.

Botanical insecticides are expected to be potential alternatives for conventional synthetic insecticides (Abdellaoui *et al.*, 2013 & Ben Hamouda *et al.*, 2015). Botanical insecticides such as agricultural waste products and plant extracts degrade quickly in air and moisture by detoxification enzymes; this refers to less environmental persistence and lower dangers to organisms that aren't the target (Isman, 2008). Rice bran extract is one of the agricultural waste products that showed a serious effect on the desert locust *S. gregaria* life cycle (Mahdy *et al.*, 2019 & 2020).

The purpose of the current study is to investigate the histopathological changes brought about by the latent effects of lufenuron and rice bran extract in newly hatched nymphs of *Schistocerca gregaria*.

MATERIALS AND METHODS

1- Insects Colony:

Desert locust *S. gregaria* were reared at the Plant Protection Research Institute, Agricultural Research Center according to the technique described by Abbassi *et al.* (2003). They are placed in wooden cages with a layer of sand (15 cm deep) on the bottom and a humidity of 10-15% suitable for spawning. A bulb (100 watts) was set to maintain a continuous photoperiod (12 light:12 dark). The locusts were maintained at a temperature of (32±2°C). Every day, feces and dead locusts were removed and the entire cage was completely cleaned and effectively sterilized at the end of any experiment.

2- Tested Compounds:

- a- Lufenuron (10%) is a chitin synthesis inhibitor.
- b- Rice bran extract is a plant waste product, it was extracted by utilizing petroleum ether through Bakr *et al.* (2006)'s method. Then it was stored in screw-capped vials at 4° C until use.

3- Bioassays Studies:

S. gregaria fifth nymphal instar females were treated using the feeding technique with LC₅₀ concentrations of lufenuron (10%) and rice bran extract which were calculated by Mahdy *et al.* (2020) and were (95 & 14.3×10³ ppm) respectively. Samples of newly hatched nymphs were collected from normal and affected eggs resulting from these treatments to study the latent effect of these compounds on different organs and jumping legs using the histological technique.

4-Histological Studies:

The samples of head capsules, abdominal region, and hind legs of control and affected newly hatching were moved to the initial fixative after being submerged in 3% glutaraldehyde. They were post-fixed for 1 hour in 1% osmium tetroxide in 0.1M cacodylate buffer following getting fixed overnight and washed three times in sucrose-cacodylate buffer (pH 7.2), washed three times in distilled water, followed by a graded sequence of ethanol alcohol dehydration. Then specimens were infiltrated with spur's epoxy resin in a graded series of absolute alcohol-spur's resin mixtures and then embedded in freshly prepared spur's resin at 70°C for 72hrs. Samples were sectioned with glass knives on a Reichert OM-2 ultra-microtome.

RESULTS AND DISCUSSION

The present study indicated that; the sublethal dose (LC₅₀) of each lufenuron and rice bran extract had latent histopathological effects on the newly hatching of affected *S. gregaria* eggs compared with control as the following:

The Head:

Semithin sections in the head of a control group showed the brain with its 3 lobes; protocerebrum, deutocerebrum, and tritocerebrum. Each one is pair, the protocerebrum is the largest part of the brain, and it continuous laterally with the optic lobe to the compound eye (Fig.1a). Neurosecretory cells in the pars intercerebralis and pars lateralis of the protocerebrum (Fig.1b). Similar structure and other details of the brain in the desert locust and others had been described by Bharadwaj & Banerjee (1971); Kurylas *et al.* (2008) and Chapman (2013).

The compound eyes consist of many similar units of vision called ommatidia. Each ommatidium consists of a convex thickened transparent cuticle known as the corneal lens. Below the lens, there are a crystalline cone and photoreceptor cells (retinula cells) to differentiate between color and brightness, light passes through the lens and cone into a channel, known as a rhabdom, two kinds of pigment cells; the primary pigment cells, completely surround the apex of the crystalline cone, and the secondary pigment cells encircle the crystalline cone and the retinula, these pigment cells separate ommatidium from its neighbors (Fig.1c). Our results are in agreement with the structure of the locust compound eye was described by Wilson *et al.* (1978); Homberg & Paech (2002); Chapman (2013) and Nation (2015).




Fig.1 (a, b & c): The semithin section in the head capsule of newly hatched nymphs of normal *S. gregaria* eggs shows a normal histological structure of (a) brain [B] and its parts: protocerebrum [Pr], deutocerebrum [De], and tritocerebrum [Tr]; The neurosecretory cells [Ns] in the protocerebrum within pars intercerebralis [PI] and pars lateralis [PL]; and the compound eye [CE]; (b) the compound eye unit (ommatidium [Om]) consists of the corneal lens [CL], crystalline cone [CC], retinula cells [RC], a rhabdom [R] the primary pigment cells [PPC] and the secondary pigment cells [SPC] and (c) the neurosecretory cells have normal nuclei.

But in the affected group with lufenuron treatment, the brain deformation was observed, it lost its basic structures and integrity, and it becomes solid masses scattered, and no longer connected to sensory organs, as well as the neurosecretory cells lysed. The compound eyes showed an asymmetrical shape and unorganized histological constituents (F.2a). Also rice bran extract caused compression and loss in the brain morphic nature, and poorly developed compound eyes (F.2b).




Fig.2 (a & b): Semithin sections in the head capsule of newly hatched nymphs produced by affected *S. gregaria* eggs with lufenuron (a) and rice bran extract (b) treatments showing the latent histopathological alteration in the brain [B] and compound eyes [CE].

These observations are in agreement with the findings of the brain and compound eyes damage in newly hatched nymphs of *S. gregaria* due to lufenuron treatment reported by (Abdel Rahman, 2017), also, rice bran extract and lufenuron exhibited disruption in the histological formation of the brain and compound eyes during embryonic development in the desert locust (Mahdy *et al.*, 2020). Similar observations showed by different plant extracts as azadirachtin on *S. gregaria* eggs caused poorly developed brain and compound eyes of newly hatching (Ghazawy *et al.*, 2010), abnormal brain cells of desert locust nymphs treated with *Cestrum* leaf powder were observed by Ben Hamouda *et al.* (2011).

The Midgut:

The midgut is the principal site for digestive enzyme production, food digestion, and nutrient absorption.

In the control group, the midgut is lined by normal enteric epithelium which has two main types of cells, the columnar epithelial cells, and regenerative cells, or nidi cells at the base of the midgut epithelium to renew destroyed epithelial cells. These cells cling to the basal lamina, continuing serially from the base to the epithelium apex. An inner layer of circular muscle and an outer layer of longitudinal muscle surrounds the midgut epithelium on the outside. There are vesicles filled with secretion at the ends of epithelial cells. The midgut epithelium is separated from the gut lumen by the peritrophic membrane (Fig.3). Similar structure was described by Chapman (2013), and Nation (2015) in the same insect and others.




Fig.3: The semithin section in the midgut of newly hatched nymphs of normal *S. gregaria* eggs shows a normal histological structure of epithelial cells [EC], regenerative cells [RC], the basal lamina [BL], circular muscle layer [CM], longitudinal muscle layer [LM], vesicles [Ve], the gut lumen [L] and peritrophic membrane [P].

Our observation showed the latent effects of each lufenuron and rice bran extract on the midgut of newly hatching, where lufenuron caused shrinkage in the gut epithelium cells which lost its normal shape with vacuoles, rupture of the circular muscle layer, and the peritrophic membrane, also vesicles spread in the gut lumen leading to blockage of the midgut (Fig.4a).

Severe histopathological effects on midgut cells caused by rice bran extract where the gut's epithelial cells lost their shape and structure leading to stretching, and deterioration of cells, as well as the vanishing of peritrophic membrane and regenerative cells. Cell content fragmentation was observed in the midgut lumen, also circular and longitudinal muscles are obliterated (Fig.4b).




Fig.4 (a & b): Semithin sections in the midgut of newly hatched nymphs produced by affected *S. gregaria* eggs with lufenuron (a) and rice bran extract (b) showing the latent histopathological alteration. [Ve] vesicles.

Similar histopathological observations were reported by many authors by using other IGRs and plant extracts as well as Bouaichi & Chihrane (2001) showed that

diflubenzuron effect on the peritrophic membrane structure in the midgut of *S. gregaria*, Tebufenozide treatments resulted in abnormal mid-gut epithelial cells of *S. gregaria* (Ghoneim *et al.*, 2008), necrosis of epithelial cells and many vacuoles resulted after treatment of *S. gregaria* and *L. migratoria* adults with azadirachtin (Nasiruddin & Mordue, 1993). The midgut of desert locusts exhibited a decrease in epithelial cell stature and muscular layer thickness after ingesting an artificial diet that included *Cestrum parquii* leaf powder (Ammar & N'cir, 2008).

These histopathological alternations agree with the observation of other species in the same order such as grasshopper *Heteracris annulosa* midgut caused by neem seed extract (Naqvi *et al.*, 1994), Also alterations of the gut structure of the grasshopper *Heteracris littoralis* reported after treatment with the *Allium sativum* essential oils (Sharaby *et al.*, 2012). On the other hand, lufenuron exhibited histopathological effects on other species; it caused the destruction of the epithelial cells of *S. littoralis* larvae's gut (Adel, 2012).

The Muscles of The Legs:

The striated muscle is the only type of muscle in insects. Histological study of the muscles in the femur of the jumping legs in newly hatched nymphs of control eggs showed that skeletal muscle tissue is formed of muscle fibers in bundles (fascicles) which are separated by a slightly thick connective tissue layer called perimysium, another thin layer of connective tissue called endomysium surrounds each muscle fiber which acting as a site of metabolic exchange, each muscle fiber has peripheral nuclei and surrounds by the plasma membrane (sarcolemma), capillaries and the fibroblasts appeared in the space between muscle fibers that maintains muscle structure (Fig.5). Similar basic muscle structure showed by Chapman (2013) and Nation (2015).




Fig.5: The semithin section in the muscles of the jumping leg femur in newly hatched nymphs of control *S. gregaria* eggs shows a normal structure of muscle tissue, muscle fiber bundles (fascicles [F]), muscle fiber [MF], perimysium [Pm], endomysium [Em], peripheral nuclei [N], sarcolemma [Sl], capillaries [C] and the fibroblasts [fb].

Histopathological examination of the leg muscles of newly hatching affected by lufenuron showed degeneration of connective tissue layers perimysium and endomysium resulting in the attachment of muscle fiber bundles, sarcolemma disappearance of some fibers, the vacuolated nuclei appears, and vanishment of other components as compared with the control muscles (Fig.6a). In affected legs by rice bran extract, complete decomposition of some fascicles and shrinkage in others occurred, the sarcolemma of some fibers, endomysium in the most bundles and some components disappeared. (Fig.6b).




Fig.6 (a & b): Semithin sections in the muscles of the jumping leg femur in newly hatched nymphs produced by affected *S. gregaria* eggs with lufenuron (a) and rice bran extract (b) showing the latent histopathological alterations.[*]degenerated muscle bundles.

Our results agree with Hussein *et al.* (2007) who showed that Rice bran extract caused histopathological effects on the muscles of the desert locust, also Bakr *et al.* (2008) and Al-Zeeb *et al.* (2018) reported that Lufenuron treatment caused malformation and degradation in the desert locust muscles. On the other hand, other IGRs as tebufenozide and Flufenoxuron Indicated disrupture in the muscle structure of the desert locust (Ghoneim *et al.*, 2008 and Bakr *et al.*, 2008).

The present study revealed that The tested compounds have latent histopathological effects where they induced serious changes in newly hatching of affected *S. gregaria* eggs with lufenuron and rice bran extract treatments and these associated with their toxic effects which were reported by Mahdy *et al.* (2019 & 2020), where they caused a disturbance in the protein and DNA of eggs resulting from treated females as fifth nymphal instars with LC₅₀ of them leading to histopathological variations of embryonic structure and inhibition of embryonic development and hatching in some eggs, and this explains, why some affected eggs produce abnormal newly hatched nymphs.

A hormonal imbalance especially juvenile hormone (JH) in affected adults female may be the main reason for all these problems where many authors explain that vitellogenesis and oocyte maturation is stimulated by Juvenile hormone (JH) as Wyatt & Davey (1996); Wu *et al.* (2021) and Zheng *et al.* (2022), also Luo *et al.* (2017) reported that vitellogenesis and egg formation are reportedly dependent on the 78-kDa glucose-regulated protein (Grp78) genes that JH regulates.

Conclusion:

Finally, our results suggest that lufenuron and rice bran extract are potential compounds for safe control against *S. gregaria* in integrated pest management (IPM).

Ethical Approval:

The study was approved by the ethical committee of the Institutional Animal Care and Use Committee, Faculty of Science, Ain Shams University (ASU-SCI/ENT/2023/5/4).

REFERENCES

- Abbassi, K.; Atay-Kadiri, Z. and Ghaout, S. (2003): Biological effects of alkaloids extracted from Three plants of Moroccan arid areas on the desert locust. *Journal Physiological Entomology*, 28: 232-236.
- Abdel Rahman, Kh. M. (2017): Embryonic development disrupted in the desert locust

- Schistocerca gregaria* Forskal (Orthoptera: Acrididae) due to lufenuron application. *Efflatounia*, 17: 1-8.
- Abdellaoui, K.; Ben Halima-Kamel, M.; Acheuk, F.; Soltani, N.; Aribi, N. and Ben Hamouda, M. H. (2013): Biochemical and histological effects of gibberellic acid on *Locusta migratoria migratoria* fifth instar larvae. *Pesticide Biochemistry and Physiology*, 107:32–37.
- Adel, M. M. (2012): Lufenuron impair the chitin synthesis and development of *Spodoptera littoralis* bosid. (Lepidoptera: Noctuidae). *Journal of Applied Sciences Research*, 8(5): 2766-2775.
- Al-Zeeb F. A. M.; Ibrahim E. H. and Bakr, R. F. A. (2018): Histopathological changes in the muscle of the desert locust, *Schistocerca gregaria* (Orthoptera: Acrididae) treated with insect growth regulator (IGR), lufenuron. *Egyptian Academic Journal of Biological Sciences (D-Histology and Histochemistry)*, 10(1): 13- 25
- Ammar, M. and N'cir, S. (2008). Incorporation of *Cestrum parquii* (Solanaceae) leaves in an artificial diet affected larval longevity and gut structure of the desert locust *Schistocerca gregaria*. *Tunisian Journal of Plant Protection*, 3(1): 27 – 34.
- Azam, K. M. and Seegh, A. A. A. (1993): Effect of diflubenzuron on second instar nymphs of desert locust, *Schistocerca gregaria* Forsk. *Journal of Research APAU / Andhra Pradesh Agricultural University*, 21(1/2): 48-50.
- Bakr, R. F. A.; Hussein, M. A.; Hamouda, L. S.; Hassan, H. A. and Elsokary, Z. F. (2008): Effect of some insecticidal agents on some biological aspects and protein patterns of desert locust *Schistocerca gregaria* (Forsk.). *Journal of Egyptian Academic Society for Environmental Development*, 9 (2): 29-42.
- Bakr, R. F.; El Bermawy, S. M.; Geneidy, N. A. M.; Emara, S. A. and Hassan, H. A. (2006): Occurrence of the biological effects of some plant extracts on the cotton leaf worm *Spodoptera littoralis* (Boisd.) and their physiological impact. *Journal of Egyptian Academic Society for Environmental Development*, 7 (1):109-147.
- Ben Hamouda, A.; Boussadia, O.; Bedis, K.; Laarif, A. and Braham, M. (2015): Studies on insecticidal and deterrent effects of olive leaf extracts on *Myzus persicae* and *Phthorimaea operculella*. *Journal of Entomology and Zoology Studies*, 3 (6): 294 -297.
- Ben Hamouda, A.; Mohamed, A. and Ben Hamouda, M. H. (2011): Effect of *Olea europaea* and *Cestrum parqui* leaves on the cuticle and brain of the desert locust *Schistocerca gregaria* Forsk. (Orthoptera: Acrididae). *Pest Technology*, 5 (1): 55- 58).
- Bharadwaj, R. K. and Banerjee, S. K. (1971): The nervous system of the desert locust, *Schistocerca gregaria* (Orthoptera: Acrididae) with a discussion on muscle innervation. *Journal of Natural History*, 5(2): 183-208.
- Bouaichi, A. and Chihrane, J. (2001): The potential and extension effect of diflubenzuron (IGR's) on the reproduction and feeding behavior of Desert Locust *Schistocerca gregaria* (Orthoptera: Acrididae), pp. 83. In: Metaleptea, Special Meeting Issue, International Conference on Orthopteroid Insects, Montpellier, France.
- Chapman, R. F. (2013): The insects: Structure and Function. 5th Edn. Cambridge, M. A. Cambridge University Press, United Kingdom.
- Coppen, G. D. A. and Jepson, P. C. (1996): Comparative laboratory evaluation of the acute and chronic toxicology of diflubenzuron, hexaflumuron and teflubenzuron against II instar desert locust, *Schistocerca gregaria* (Orthopetra: Acrididae). *Pesticide Science*, 46 (2): 183-190.
- Fonseca, A. P. P. ; Marques, E. J. ; Torres, J. B.; Silva, L. M. and Siqueira, H. A. A. (2015): Lethal and sublethal effects of lufenuron on sugarcane borer *Diatraea flavipennella* and its parasitoid *Cotesia flavipes*. *Ecotoxicology*, 24(9):1869–1879.

- Garriga, M. and Caballero, J. (2011): Insights into the structure of urea-like compounds as inhibitors of the juvenile hormone epoxide hydrolase (JHEH) of the tobacco hornworm *Manduca sexta*: analysis of the binding modes and structure-activity relationships of the inhibitors by docking and CoMFA calculations. *Chemosphere*, 82(11): 1604-1613.
- Ghazawy, N. A.; Awad, H. H.; Abdel Rahman, K. M. (2010): Effects of azadirachtin on embryological development of the desert locust *Schistocerca gregaria* Forskal (Orthoptera: Acrididae). *Journal of Orthoptera Research*, 19(2): 327-332.
- Ghoneim, K. S.; Bakr, R. F.; Tanani, M. A.; Al Dali, A. G. and Bream, A. S. (2008): An ultrastructural study on the desert locust (*Schistocerca gregaria*) as affected by Tebufenozide (RH-5992). *International Journal of Agriculture and Biology*, 10: 479–80.
- Homberg, U. (2015): Sky compass orientation in desert locusts-evidence from field and laboratory studies. *Frontiers in Behavioral Neuroscience*, 9:346.
- Homberg, U. and Paech, A. (2002): Ultrastructure and orientation of ommatidia in the dorsal rim area of the locust compound eye. *Arthropod Structure & Development*, 30(4):271–280.
- Hussein, M. A.; Elsokary, Z. F.; Bakr, R. F. A.; Hamouda, L. S. and Hassan H.A. (2007): Toxicity and ultrastructural studies of thoracic muscles of *Schistocerca gregaria* (Forsk.) treated with some insecticidal agents. *Journal Egyptian Academic Society For Environment Development, (A-Entomology)*, 8(3):45-54.
- Isman, M. B. (2008): Botanical insecticides: for richer, for poorer. *Pest Management Science*, 64(1): 8-11.
- Kurylas, A. E.; Rohlfing, T.; Kroficzik, S.; Jenett, A. and Homberg, U. (2008): Standardized atlas of the brain of the desert locust, *Schistocerca gregaria*. *Cell Tissue Research*, 333(1):125–145.
- Lecoq, M. (2001): Recent progress in desert and migratory locust management in Africa. Are preventive actions possible? *Journal of Orthoptera Research*, 10: 277-291.
- Luo, M.; Li, D.; Wang, Z.; Guo, W.; Kang, L. and Zhou, S. (2017): Juvenile hormone differentially regulates two Grp78 genes encoding protein chaperones required for insect fat body cell homeostasis and vitellogenesis. *Journal of Biological Chemistry*, 292(21): 8823–34.
- Mahdy, N. M.; Ahmed, S. S.; Mohamed, M.; Badawy, N. and Abdou, M. A. (2020): Histopathological and ultrastructure changes in the embryonic development of *Schistocerca gregaria* (Forsk.) (Orthoptera): (Acrididae) Induced by lufenuron (CSI) and rice bran extract (waste product). *The Egyptian Journal of Histology*, 43(1):186-203.
- Mahdy, N. M.; Mohamed, M. I.; Abdou, M. A.; Ahmed, S. S. and Badawy, N. S. (2019): Toxicological and biochemical effects of lufenuron and rice bran on desert locust, *Schistocerca gregaria* (Forsk.) (Orthoptera: Acrididae). *Egyptian Academic Journal of Biological Sciences (F.Toxicology and Pest Control)*, 11(2): 37– 58.
- Naqvi, S. N.; Tabassum, R.; Azmi, M. A.; hafez, A.; Tariq, R. M. and Rashid, N. (1994): Histopathological effects of danitol (Fenprothrin) and neem fraction on grasshopper, *Heteracris annulosa* (Wak) gut and changes in enzyme pattern. *Proceedings of Pakistan Congress of Zoology*, 14: 2532.
- Nasiruddin, M. and Mordue, A. J. (1993). The effect of azadirachtin on the midgut histology of the locust, *Schistocerca gregaria* and *Locust migratoria*. *Tissue and Cell*, 25(6): 875-884.
- Nation, J. L. (2015): *Insect Physiology and Biochemistry*. 3rd Edition, CRC Press, Boca Raton London, New York, Washington.

- Said, S. M. (2020): Histopathological, biological and biochemical effects of lufenuron on *Schistocerca gregaria* (Orthoptera: Acrididae). *Egyptian Journal of Plant Protection Research*, 3 (4): 1151-1161.
- Sampson, B. J.; Marshall, D. A.; Smith, B. J.; Stringer, S. J.; Werle, C. T.; Magee, D. J. and Adamczyk, J. J. (2017): Erythritol and Lufenuron detrimentally alter age structure of wild *Drosophila suzukii* (Diptera: Drosophilidae) populations in blueberry and blackberry. *Journal of Economic Entomology*, 110 (2):530-534.
- Sharaby, A.; Montasser, S. A.; Mahmoud, Y. A. and Ibrahim, S. A. (2012): Natural plant essential oils for controlling the grasshopper (*Heteracris littoralis*) and their pathological effects on the alimentary canal. *Ecologia Balkanica*, 4: 39–52
- Simpson, S. J.; Mccaffery, A. R. and Hägele, B. F. (1999): A behavioural analysis of phase change in the desert locust. *Biological Review of the Cambridge Philosophical Society*, 74:461-480.
- Taha, G. Z. and El-Gammal, A. M. (1990): Morphogenetic effects of non-terpenoid juvenile hormone analogue, S-31183 on, metamorphosis of last nymphal instar of *Schistocerca gregaria*. *Egyptian Journal of Applied Sciences*, 5: 75-81.
- Verloop, A. and Ferrell, C. D. (1977): Benzoylphenyl ureas-a new group of larvicides interfering with chitin deposition. In 'Pesticide Chemistry in the 20th Century'. (Ed. J. R. Plimmer). *American Chemical Society Symposium Series*, 37: 237- 270.
- Wilson, M.; Garrad, P. and McGinness, S. (1978): The unit structure of the locust compound eye. *Cell and Tissue Research*, 159: 205-226.
- Wu, Z.; Yang, L.; He, Q. and Zhou, S. (2021): Regulatory mechanisms of vitellogenesis in Insects. *Frontiers in Cell and Developmental Biology*, 8: 1863.
- Wyatt, G. R. and Davey, K. G. (1996): Cellular and molecular actions of juvenile hormone. II. Roles of juvenile hormone in adult insects. *Advances in insect physiology*, (26): 2 –155.
- Zheng, H.; Wang, N.; Yun, J.; Xu, H.; Yang, J. and Zhou, S. (2022): Juvenile hormone promotes paracellular transport of yolk proteins via remodeling zonula adherens at tricellular junctions in the follicular epithelium. *PLOS Genetics*, 18(6): e1010292.

ARABIC SUMMARY

التأثيرات الهستوباثولوجية الكامنة لمركب الليوفنيرون ومستخلص نخالة الأرز على حشرة الجراد الصحراوي (شيسستوسيركا جريجاريا)

نوره محمد مهدي¹ سميرة محمد نبيل عبد الواحد¹ أميرة السيد عبد الحميد²
¹ - معهد بحوث وقاية النباتات- مركز البحوث الزراعيّة- الجيزة - مصر
² - قسم علم الحشرات - كلية العلوم - جامعة عين شمس- القاهرة - مصر

تم التقييم المعملّي للتأثيرات الهستولوجية الممتدة (الكامنة) لكل من الليوفنيرون ومستخلص نخالة الأرز على حوريات الفقس حديثًا للجراد الصحراوي الناتجة من اناث عوملت في العمر الخامس بالجرعة النصف مميتة لكلا المركبين. واثبتت الدراسة حدوث تشوهات نسيجية كبيرة في بنية الدماغ، والعين المركبة، والمعوي المتوسط، وعضلات الساق الخلفية للفقس مقارنة بالكنترول، وفقدت هذه الأعضاء تركيبها وشكلها ووظيفتها الطبيعية. فكان للمركبين تأثيرات هستوباثولوجية خطيرة طويلة المدى، لذا من الممكن استخدامهما للسيطرة على حشرة الجراد الصحراوي كعوامل بديلة آمنة لمبيدات الآفات الاصطناعية السامة.

الكلمات المفتاحية:

الجراد الصحراوي- دراسة هستولوجية - الليوفنيرون - مستخلص نخالة الأرز - مثبطات تكوين الكيتين- منتجات المخلفات الزراعية.